

Antimicrobial activity of *Melissa officinalis* L. and *Crocus sativus* L. against oral pathogens: Detection of cellular structural changes by FT-IR.

Anastasaki E.¹, Zoumpopoulou G.², Papadimitriou K.², Tarantilis P.¹, Polissiou M.¹ and Tsakalidou E.²

¹Laboratory of Chemistry, Department of Science, Agricultural University of Athens, 75 Iera Odos Street, 11855, Athens, Greece
²Laboratory of Dairy Research, Department of Food Science and Technology, Agricultural University of Athens, 75 Iera Odos Street, 11855, Athens, Greece

Periodontal diseases and dental caries are common oral disorders in human population with a multifactorial etiology closely related with the development of dental plaque. The latter is composed of native oral microbiota and it is accumulated on teeth surfaces. Several antiseptic agents are used widely to inhibit bacterial growth [1,2]. However, these substances have adverse effects. In the current study, *Melissa officinalis* L. and *Crocus sativus* L. extracts were tested as potential natural antimicrobial agents. The antimicrobial activity of plants extracts was studied towards Gram-positive strains belonging to *Streptococcus* species related to the oral health. Fourier transform infrared spectroscopy (FT-IR) was applied in order to evaluate the changes in the cellular composition of target bacterial cells after their exposure to extracts of both plants.

Sample preparation. Plants were subjected to sequential extraction with petroleum ether, hexane, diethyl ether and methanol, as shown in figure 1.

All extracts were evaporated under reduced pressure and dried using rotary evaporator. Dried extracts were stored in labeled screw capped bottles at -20°C.




Figure 1. Schematic plan of extraction procedure

Screening of plants extracts against oral pathogens. Different concentrations of extracts were tested against six *Streptococcus* strains by the well diffusion assay (WDA).

Diethyl ether (DE) and methanol extracts (ME) of both plants induced the highest bactericidal effect against all tested bacteria, followed by petroleum ether (PE) and hexane extracts (HE) (Table 1).

Based on these results the methanolic extracts of both plants were selected for further investigation against three *Streptococcus* strains, namely *Str. mutans* LMG 14558^T, *Str. oralis* LMG 14532^T and *Str. sobrinus* LMG 14641^T.

Table 1. Antimicrobial activity of *Melissa officinalis* L. and *Crocus sativus* L. extracts towards six *Streptococcus* strains as determined by the well diffusion assay

Strain	Inhibition of <i>Melissa officinalis</i> L. (diameter; mm)									
	PE (mg/mL)		HE (mg/mL)		DE (mg/mL)		ME (mg/mL)			
	50	25	50	25	50	25	280	168	84	28
<i>Streptococcus gordonii</i> LMG 14518 ^T	11	8	ND*	ND	16	14	17	15	12	7
<i>Streptococcus mutans</i> LMG 14558 ^T	7	0	7	0	17	12	9	7	0	0
<i>Streptococcus oralis</i> LMG 14532 ^T	11	7	11	ND	26	21	14	13	12	9
<i>Streptococcus salivarius</i> LMG 11489 ^T	7	0	ND	ND	12+8	11+6	13	12	10	9
<i>Streptococcus sanguinis</i> DSM 20068	10	0	10	ND	17	14	15	14	12	9
<i>Streptococcus sobrinus</i> LMG 14641 ^T	11	9	ND	8	13	10	10	10	7	0

Strain	Inhibition of <i>Crocus sativus</i> L. (diameter; mm)									
	PE (mg/mL)		HE (mg/mL)		DE (mg/mL)		ME (mg/mL)			
	50	25	50	25	50	25	280	168	84	28
<i>Streptococcus gordonii</i> LMG 14518 ^T	8	8	ND	0	27	20	20	18	16	12
<i>Streptococcus mutans</i> LMG 14558 ^T	15	12	ND	0	40	25	16	15	13	7
<i>Streptococcus oralis</i> LMG 14532 ^T	10	8	ND	0	27	22	17	16	15	13
<i>Streptococcus salivarius</i> LMG 11489 ^T	0	0	ND	0	28	20	15	15	11	8
<i>Streptococcus sanguinis</i> DSM 20068	0	0	ND	0	30	23	20	19	18	13
<i>Streptococcus sobrinus</i> LMG 14641 ^T	8	8	ND	8	25	25	20	18	16	14

*ND= Not done

Time killing studies of methanolic extracts against *Str. mutans* LMG 14558^T, *Str. oralis* LMG 14532^T and *Str. sobrinus* LMG 14641^T. Antimicrobial activity was studied (*in vitro* killing assays) against target cells in the logarithmic phase of bacterial growth.

The viability of *Streptococcus* cells was studied for 24 hours of incubation with 28 mg/mL (final concentration) methanolic plants extracts. 99% cell death of *Str. sobrinus* LMG 14641^T was achieved in 4 hours, while for *Str. mutans* LMG 14558^T and *Str. oralis* LMG 14532^T within 6 hours for both plants extracts (Figure 2).






Figure 2. Time killing studies of logarithmic cells of *Str. mutans* LMG 14558, *Str. oralis* LMG 14532 and *Str. sobrinus* LMG 14641 after their incubation with the control sample (MeOH 70%) (●); 28 mg/mL methanolic extract of *Crocus sativus* L. (●); 28 mg/mL methanolic extract of *Melissa officinalis* L. (▲).

FT-IR Analysis. Fourier transform infrared spectroscopy (FT-IR) was applied in the respective time period, where 99% of cell death was achieved, in order to evaluate the changes in the cellular composition of cells.

The FTIR spectrum of a biological system like bacteria is complex and consists of broad bands (Figure 3) that arise from the superposition of absorption peaks of various contributing macromolecules (proteins, lipids, polysaccharides, and nucleic acids) [3]. The FT-IR spectra of control cells were compared with the spectra of incubated with methanolic extracts cells in four different regions:

Region I- 3000– 2800 cm^{-1} related to CH from fatty acids of the bacterial cell membrane

Region II- 1800– 1500 cm^{-1} related to C=O and N–H from proteins

Region III- 1500– 1200 cm^{-1} related to PO_2 from nucleic acids, as well as proteins and fatty acids

Region IV- 1200– 900 cm^{-1} : related to various absorptions of polysaccharides of the cell wall




Figure 3. FT-IR spectra of *Str. oralis* LMG 14532^T cells and the characteristic regions used for comparison

Principal component analysis (PCA) of the second derivative transformed spectra was performed for each characteristic spectral region (Figure 4) [4].

PCA revealed structural changes among cells treated with the extracts or the control sample. The significant differences were observed in characteristic spectral regions correlated to the above cellular structural components.




Figure 4. PCA of second derivative transformed FT-IR spectra of *Str. mutans* LMG 14558 (A), *Str. oralis* LMG 14532 (B) and *Str. sobrinus* LMG 14641 (C) cells in the four characteristic spectral regions after their incubation with the control sample (MeOH 70%) (●); 28 mg/mL methanolic extract of *Crocus sativus* L. (●); 28 mg/mL methanolic extract of *Melissa officinalis* L. (▲).

Conclusions. The results have shown that *Melissa officinalis* L. and *Crocus sativus* L. extracts consist of important secondary metabolites in the search for new effective antibacterial agents against the pathogens responsible for dental caries.

Diethyl ether and methanol extracts were more potent than petroleum ether and hexane extracts and were found to be prominently active.

FT-IR analysis along with chemometric analysis (PCA) of incubated *Streptococcus* cells revealed significant differences in all regions of spectra that correspond to cellular structural components.

[1] Marsh, P. D. (2006). Dental plaque as a biofilm and a microbial community: Implications for health and disease. BMC Oral Health, 6, 214.

[2] Jenkinson, H. F., Lala, H. C., & Shepherd, M. G. (1990). Coaggregation of *Streptococcus sanguis* and other streptococci with *Candida albicans*. Infection and Immunity, 58, 1429–1436.

[3] Kansiz, M., Heraud, P., Wood, B., Burden, F., Beardall, J., McNaughton, D., (1999). Fourier transform infrared microspectroscopy and chemometrics as a tool for the discrimination of cyanobacterial strains. Phytochemistry 52, 407–417.

[4] Papadimitriou, K., Boutou, E., Zoumpopoulou, G., Tarantilis, P.A., Polissiou, M., Vorigas, C.E., Tsakalidou, E., (2008). RNA arbitrarily primed PCR and Fourier transform infrared spectroscopy reveal plasticity in the acid tolerance response of *Streptococcus macedonicus*. Applied and Environmental Microbiology 74, 6068–6076.



This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program: Heracleitus II. Investing in knowledge society through the European Social Fund.

